

Ladasten® instructions

translated from original Russian instructions by Extrapharmacy Online Store

<http://extrapharmacy.ru>

Active substance : Adamanthybromphenylamin

Pharmacologic action : anti-asthenic

Ladasten is a derivative of Adamantane. It impacts positive on physical and mental performance. The spectrum of action of Ladasten is a combination of activating, anxiolytic, immunostimulant actions and actoprotective activity. Ladasten doesn't have any hypno-sedative or muscle relaxant properties, not addictive.

Therapeutic effect of Ladasten in patients with asthenia and anxiety-asthenic disorders is manifested from the first day of its application . Ladasten helps to restore the activity and increase endurance.

Ladasten is nontoxic

Pharmacokinetics

The time to maximum concentration (Tmax) is about 2-4 hours,
maximum concentration (Cmax) is 363.3 ng /ml,
half-life period (T1 / 2) is 11.21 hours

Indications

Asthenic conditions of various origins, including somatic diseases and after infectious diseases.
Neurasthenia.

Contraindications

Pregnancy, lactation, children under 18 years , hypersensitivity to the drug.

Side effects

Manifestations of excessive activation and sleep disorders are possible, which do not require discontinuation of the drug, but it is advisable to decrease the dose. With increased individual sensitivity to the drug - allergic reactions may develop.

Interaction

Ladasten reduces the hypnotic effect of thiopental sodium, does not weaken the anxiolytic effect of benzodiazepines.

Dosing and Administration

Ladasten is taken orally regardless of the meal. The optimal single dose - 50-100 mg; daily dose - 100-200 mg, distributed into 2 doses throughout the day.

The drug should not be used after 4 pm. The duration of a course of Ladasten is 2-4 weeks.

Overdose

With substantial overdose - sedation may develop. Treatment: nonspecific detoxification therapy.

Manufacturer : LEKKO, Russia

Reliable supplier with fast Worldwide shipping

Extrapharmacy Online Store

<http://extrapharmacy.ru>

Storage

The temperature is not above 25 ° C.

Keep out of the reach of children.

Shelf-life of the drug is 2 years.