

# Solcoseryl solution for injections 42.5mg/ml

translated from original Russian instructions by Extrapharmacy Online Store

<http://extrapharmacy.ru>

Name in Cyrillic : СОЛКОСЕРИЛ

**Active ingredient:** deproteinized dialysate from the blood of healthy dairy calves.

## Indications:

Disorders of the peripheral arterial and venous circulation: occlusive peripheral arterial disease in stages III-IV(Fontaine); chronic venous insufficiency, accompanied by trophic disorders.

Violations of cerebral metabolism and blood circulation: ischemic stroke; hemorrhagic stroke; traumatic brain injury.

## Contraindications:

Children and teens under 18 years of age (safety data applications are not available); pregnancy (use of safety data are not available); lactation (safety data applications are not available); established hypersensitivity to bovine blood dialysate; Hypersensitivity to derivatives of para-hydroxybenzoic acid (E216 and E218) and the free benzoic acid (E210).

Use with care for patients with hyperkalemia, renal failure, cardiac arrhythmias, with concomitant drugs potassium (as Solkoseril contains potassium), with oliguria, anuria, pulmonary edema, severe heart failure.

## Dosing and Administration:

The drug is administered intravenously (drip) (previously diluted with 250 mL of 0.9% sodium chloride or 5% dextrose solution, sodium), slow intravenous injection (pre-diluted with 0.9% sodium chloride or 5% dextrose solution, sodium in a ratio of 1: 1) or intramuscularly

Peripheral arterial occlusive disease in stages III-IV on Fontaine: intravenously daily with 20 ml. Duration of therapy up to 4 weeks and is determined by the clinical picture of the disease.

Chronic venous insufficiency, accompanied by trophic disorders: intravenously 10 ml, 3 times a week. Duration of therapy is no more than 4 weeks and is determined by the clinical picture of the disease.

If you have a local trophic disorders of tissues the therapy with Solkoseryl-gel and then Solkoseryl-ointment is recommended.

Traumatic brain injury, metabolic and vascular diseases of the brain: 10-20 ml intravenously daily for 10 days. Next - intramuscular or intravenous injection of 2 ml to 30 days.

If intravenous administration is not possible, the drug can be administered by intramuscular injection of 2 ml / day

## Cautions:

Interactions: Be careful to apply simultaneously with drugs that increase blood potassium (potassium supplements, potassium-sparing diuretics, ACE inhibitors).

The drug should not be confused when administered with other drugs (especially phytoextracts).

The preparation of parenteral forms is incompatible with Ginkgo biloba, and naftidrofuryl and bentsiklana fumarate.

Side effects: Allergic reactions: seldom - rash, fever.

Local reactions: rarely - redness, swelling at the injection site.

Special instructions: should stop using the product and Symptomatic treatment In case of adverse reactions.

**Manufacturer** Valeant Pharma, Switzerland

**Reliable supplier with fast Worldwide shipping**

Extrapharmacy Online Store

<http://extrapharmacy.ru>

**Storage** : Store in dark place, the temperature is not higher than 25° C. Shelf-life of the drug is 5 years.